

PLASTIC EYE SURGERY ASSOCIATES, PLLC

James R. Patrinely, MD, FACS | Charles N.S. Soparkar, MD, PHD, FACS

3730 Kirby Drive, Suite 900, Houston, TX 77098

Telephone (713) 795-0705 / FAX (713) 807-0630

THE WEEKS BEFORE SURGERY:

To minimize the risk for bleeding during surgery, bruising afterwards, and possible compromise of your surgical outcome, it is important that you review the following lists.

These aspirin-like medicines should be stopped three weeks prior to your surgery. Please review the list carefully.

If you are taking any of these medicines on the advice or direction of a physician, you should consult that doctor before making any changes.

© PLASTIC EYE SURGERY ASSOCIATES, PLLC

03/06

Acetylsalicylic Acid	Buffaprin	Fiorinal	Salicylic Acid
Alka-Seltzer	Bufferin	Magnaprin	Triaminicin
Anacin	Buffex	Measurin	Trigesic
Arthritis Pain Relievers	Buffinol	Midol	Ursinus
ASA	Ecotrinol	Momentum	Vanquish
Ascriptin	Empirin	Norgesic	Zorprin
Aspergum	Excedrin	Panwarfin	
Aspirin	4-Way Cold Tablets	Plavix	

The following medicines need to be stopped at least 2 weeks prior to surgery:

Advil	Dolobid	Meclomen	Rufen
Anaprox	Feldene	Medipren	Salsalate
Ansaid	Fenoprofen	Monogesic	Salsitab
Anturane	Flurbiprofen	Motrin	Sulindac
Butazolidin	Genpril	Nalfon	Tolectin
Celebrex	Haltran	Naprosyn	Tolmetin
Clinoril	Heparin	Naproxen	Tordol
Coumadin	Ibuprin	Nuprin	Trendar
Diclofenac	Ibuprofen	Orudis	Vioxx
Dicumarol	Indameth	Pamprin	Voltaren
Diflunisal	Indocin	Persantine	Warfarin
Dipyridamole	Indomethacin	Phenylbutazone	
Disalcid	Ketoprofen	Piroxicam	
	Meclodium	Ponsteel	

The following supplements should be stopped 2 weeks before surgery:

Garlic	Ginseng
Ginger	Vitamin E
Ginko Biloba	